


Post Copy

POV: Your filler was designed with expression in mind.


The Teoxane RHA® Collection | Filler designed to adapt to your face for natural-looking results.

Contact us to book your RHA® appointment today!

#TeoxaneRHACollection #RHACollection
#JoinTheMovement



SLIDE 1



SLIDE 2

SLIDE 3 SCROLLING ISI

The Teoxane RHA® Collection includes hyaluronic acid (HA) fillers with lidocaine and mepivacaine. Revance will only distribute the Teoxane RHA® Collection with mepivacaine, thus may refer to the products without the use of "Mepi" in the product name. The long-term efficacy and safety outcomes of RHA® with lidocaine are applicable to RHA® with mepivacaine, and mepivacaine has shown no significant impact on gel properties, including rheology and degradation profile.

RHA® Collection of Fillers, by Teoxane

Approved Uses

The Teoxane RHA® Collection of resilient hyaluronic acid (HA) fillers includes RHA Redensity®, RHA® 2, RHA® 3 and RHA® 4 with lidocaine and RHA Redensity® Mepi, RHA® 2 Mepi, RHA® 3 Mepi and RHA® Dynamic Volume, with mepivacaine.

RHA Redensity® and RHA Redensity® Mepi are for injection into the facial tissue for the correction of moderate to severe dynamic perioral rhytids. RHA® 2, RHA® 2 Mepi and RHA® 4 are for the correction of moderate to severe dynamic facial wrinkles and folds, such as nasolabial folds (NLFs), in adults 22 or older. RHA® 3 and RHA® 3 Mepi are for the correction of moderate to severe dynamic facial wrinkles and folds, such as nasolabial folds (NLFs), and also used to augment lip fullness in adults 22 or older. RHA® Dynamic Volume is for correction of moderate to severe dynamic facial wrinkles and folds, such as nasolabial folds (NLFs), and for cheek augmentation and/or correction of age-related midface contour deficiencies in adults 22 or older.

IMPORTANT SAFETY INFORMATION

Are there any reasons why I should not receive any RHA® injectable gel formulation?

Do not receive if you have a history of multiple severe allergies or severe allergic reactions; if you are allergic to lidocaine or mepivacaine or gram-positive bacterial proteins; or if you have a bleeding disorder.

What precautions should I discuss with my doctor?

- Tell your doctor if you are pregnant or breastfeeding as the safety of these products for use during pregnancy or while breastfeeding has not been studied.